Indirect Down-regulation of Tumor-suppressive *let-7* Family MicroRNAs by *LMO1* in Neuroblastoma

NORIHISA SAEKI¹, AKIRA SAITO², YUKI SUGAYA², MITSUHIRO AMEMIYA² and HIROKI SASAKI³

¹*Division of Anatomy and Physiology, Okinawa Prefectural College of Nursing, Okinawa, Japan;*

²*Statistical Genetics Analysis Division, StaGen Co. Ltd., Tokyo, Japan;*

³*Department of Translational Oncology, National Cancer Center Research Institute, Tokyo, Japan*

Abstract. *Background/Aim:* Overall survival for the high-risk group of neuroblastoma (NB) patients still remains at 40-50%, necessitating the establishment of a curable treatment. *LIM domain only 1 (LMO1)* gene encoding a transcriptional regulator is an NB-susceptibility gene with a tumor-promoting activity. Previously we conducted chromatin immunoprecipitation and DNA sequencing analyses on NB cell lines and identified 3 protein-coding genes regulated by *LMO1*. In this study, we extended our analyses to capture microRNA genes directly or indirectly regulated by *LMO1*. *Materials and Methods:* Using microarrays, we conducted a comparative gene expression analysis on an NB cell line SK-N-SH; between the cells with and without *LMO1* suppression. *Results:* Overall, 18 microRNAs were identified to be indirectly down-regulated by *LMO1* including 7 microRNAs of the *let-7* family, whose cell proliferation inhibitory activity was observed. *Conclusion:* Target genes of the *LMO1*-regulated microRNAs and their relevant pathways may be a potential therapeutic target.

Neuroblastoma (NB), the most common extracranial solid tumor in childhood, is a cause for approximately 15% of all paediatric cancer deaths (1). With recent progress in NB treatment, patient prognosis has been significantly improved; Overall survival (OS) for low-risk and intermediate-risk groups is now >98% and 90-95%, respectively. On the contrary, OS for the high-risk (HR) group is still at 40-50%, and 50-60% of these patients have a relapse, for which no

option of curable treatment remains (2, 3). To improve the OS and quality of life for the HR group, the identification of novel therapeutic target molecules is imperative (4-6).

Previously, the LIM domain only 1 (*LMO1*) gene encoding a transcriptional regulator was identified as an NB susceptibility gene by a genome-wide association study (7). In the study, it was also revealed that both the presence of an NB-related allele of single-nucleotide polymorphism (SNP) rs110419 in the *LMO1* gene and the increased copy number of the gene contribute to augmented *LMO1* expression, and that *LMO1* promotes proliferation of NB cells. Importantly, it was shown that the relation of the variation at rs110419 in *LMO1* as well as the increased copy number to NB is stronger in the HR group than in the non-HR group. Recently, the SNP rs110419 was found to be located in an enhancer region of the gene and have an influence on its function (8). Consequently, it is likely that the transcriptional regulator *LMO1* is involved in the regulation of expression of a variety of genes important for NB progression in HR patients, and it is anticipated that identification of the *LMO1*'s regulatory targets may lead us to identify novel tumor-promoting molecular pathways in NB, as well as novel therapeutic target molecules, especially for treatment of the HR group.

Previously, we conducted chromatin immunoprecipitation and DNA sequencing analyses on two NB cell lines, SK-N-SH and LA-N-5 and identified 3 protein-coding genes regulated by *LMO1* in the 2 NB cells: LIM and senescent cell antigen-like domains 1 (*LIMS1*), Ras suppressor protein 1 (*RSU1*) and relaxin 2 (*RLN2*) (9). *LIMS1* and *RSU1* encode proteins functioning with integrin-linked kinase (ILK) in integrin adhesomes, and an ILK inhibiting compound Cpd22 inhibited proliferation of the NB cells *in vitro* (9-12). The findings suggest the possibility that in addition to the *LIMS1*/ILK pathway, more tumor-promoting pathways exist downstream of the *LMO1*-regulatory cascade.

As the previous study aiming at *LMO1*-regulated genes was limited to protein-coding genes, analyses were extended to microRNA genes in this study. MicroRNAs are known to

This article is freely accessible online.

Correspondence to: Norihisa Saeki, Division of Anatomy and Physiology, Okinawa Prefectural College of Nursing, Yogi 1-24-1, Naha, Okinawa 902-0076, Japan. Tel: +81 988338885, Fax: +81 988335133, e-mail: saeki@okinawa-nurs.ac.jp

Key Words: Neuroblastoma, pediatric cancer, *LMO1*, microRNA, *let-7*.

have an important role in regulating gene function by binding to the 3' untranslated region (UTR) of target messenger RNAs (mRNAs) and suppressing their translation (13, 14). Therefore, it is likely that the mRNAs targeted by LMO1-regulated microRNAs are significantly involved in NB oncogenesis.

Materials and Methods

Cell line. The NB cell line SK-N-SH was provided from the European Collection of Cell Cultures and maintained in D-MEM (Wako Pure Chemical Industries, Ltd., Osaka, Japan) supplemented with 10% fetal calf serum. LA-N-5 was provided by the RIKEN BRC through the National Bio-Resource Project of the MEXT, Japan, and maintained in RPMI1640 (Life Technologies Japan Ltd., Tokyo, Japan) supplemented with 10% FCS. SK-N-BE (2) was obtained from ECACC and maintained in a 1:1 mixture of EMEM (Wako) and Ham's F12 (Wako) supplemented with 0.5% non-essential amino acids (Life Technologies) and 15% FCS.

shRNA. Each shRNA construct was prepared by annealing the (a) strand to the (b) as follows: for shLMO1 (target sequence: 5'-GGCATTGGACAAGTACTGG-3'), (a) 5'-tcgagGGtATTGGAtAAGTAtTGGttcaagagaCCAGTACTTGTCCAATGCC-tttttacgcgta-3' and (b) 5'-agcttacgcgtaaaaaGGCATTGGAGAAAGTACTGGtctcttgaaCCAACTATTCCAATaCCc-3', for shGFP (target sequence: 5'-GCTACCTGTTCCATGGCCAA-3'), (a) 5'-tcgagGCTAtCTGTTCCgTGGCCgAttcaagagaTTGGCCATGGAACAGGTAGttttttacgcgtg-3' and (b) 5'-gatccacgcgtaaaaaCTACCTGTTCCATGGCCAAAtctcttgaaTcGGCCAcGGAACAGaTAGCc-3'. The shRNA was inserted into pLVSiN-CMV neo (Takara Bio, Shiga, Japan) whose CMV promoter was replaced by hU6 promoter. To obtain viral particles, the pLVSiN-hU6-shRNA constructs were introduced into Lenti-X™ 293T Cells (Takara Bio) with Lenti-X™ HTX Packaging System (Takara Bio), and after 72 h incubation, the medium was collected and its viral titer (infection units/ml) was determined by transduction to HT1080 cells. The NB cell lines were transduced with lentivirus (> 100,000 infection units) in the presence of polybrene (10 µg/ml in culture medium, Sigma-Aldrich, St. Louis, MO, USA).

Microarray expression analyses. Total RNA was extracted from SK-N-SH transduced with pLVSiN-shLMO1 or pLVSiN-shGFP, using Absolutely RNA miRNA Kit (Agilent Technologies, Santa Clara, CA, USA). Microarray expression analyses were performed with miRNA Complete Labeling and Hyb Kit (Agilent) and SurePrint G3 Human miRNA Microarray, Release 19.0 (Agilent), following a standard protocol recommended by the manufacturer. The data were analyzed using Agilent Feature Extraction Software (Agilent).

Real-time RT-PCR. For microRNA hsa-miR-3648, RT-PCR was conducted by converting about 10 ng of total RNA to the first strand cDNA with TaqMan MicroRNA Reverse Transcription Kit (Life Technologies), followed by performing TaqMan MicroRNA Assay (Life Technologies, Applied Biosystems Assay ID: 464401). For mRNA, RT-PCR was performed by converting approximately 5 µg of total RNA to the first strand cDNA with High Capacity cDNA Reverse Transcription Kit (Life Technologies), followed by conducting TaqMan Gene Expression

Table I. MicroRNA genes present in the vicinity of the peaks detected by ChIP-Seq for LMO1-targets.

Chromosome	Peak position [†]		MicroRNA gene*
	From	To	
1	84315912	84315949	<i>hsa-mir-548ap</i>
3	120131193	120132019	<i>hsa-mir-198</i>
5	134260148	134260380	<i>hsa-mir-4461</i>
5	134262667	134262849	<i>hsa-mir-4461</i>
5	134263306	134263645	<i>hsa-mir-4461</i>
21	9826946	9827631	<i>hsa-mir-3648</i> , <i>hsa-mir-3687</i>

[†]The position is based on UCSC hg19. *Present within 5kb from the peak position.

Assay (Life Technologies, Applied Biosystems Assay ID: Hs00231133_m1 for LMO1 and Pre-Developed TaqMan Assay Reagent for glyceraldehyde-3-phosphate dehydrogenase (GAPDH)). The PCR was performed for 40 cycles that included 2 steps of temperature: 95°C for 15 sec and 60°C for 60 sec, using ABI PRISM 7900HT Sequence Detection System (Life Technologies). The relative transcript level was calculated using the Ct value of GAPDH transcript as reference.

Down-regulation of GAPDH and western blot analyses. Control siRNA and siRNA for GAPDH suppression were purchased from BIONEER Corp, Daejeon, Korea, and introduced into NB cells using Lipofectamine RNAiMAX (Life Technologies). Using CellLytic-M Mammalian Cell Lysis/Extraction Reagent (Sigma-Aldrich) and Protease Inhibitor Cocktail (Sigma-Aldrich), cell lysates were prepared and loaded onto a 15% SDS-polyacrylamide gel. Immune blotting was performed with an anti-GAPDH, an anti-Actin and a HRP-conjugated anti-goat antibody (Santa Cruz Biotechnology, Dallas, Texas, USA). The signals were detected by Pierce Western Blotting Substrate Plus (Thermo Fisher Scientific, Yokohama, Japan).

Cell growth assay. A mirVana miRNA inhibitor for miR-3648 and a negative control were obtained from Life Technologies. The microRNA mimics and Negative control #2 were purchased from BIONEER Corp. The NB cells were prepared in five wells of 96-well plates (10,000 cells/well). The microRNA inhibitor and mimics were added to the wells (6 pmol/well) using Lipofectamine RNAiMAX (Life Technologies). The cells were incubated for 72 h and the growth was evaluated by MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) assay using Cell Counting Kit-8 (Dojindo, Kumamoto, Japan).

Results

MicroRNA hsa-miR-3648 is a direct target of LMO1 regulation. In our previous study, we performed chromatin immunoprecipitation-DNA sequencing (ChIP-Seq) to identify LMO1-regulated genes using 2 NB cell lines SK-

Figure 1. Down-regulation of *LMO1* by shRNA (real-time RT-PCR). The shRNA for *LMO1* (*shLMO1*) down-regulates *LMO1* expression in NB cell lines, SK-N-SH and LA-N-5, compared to the cells transfected with shRNA for green fluorescent protein (*shGFP*), statistical significance at $p < 0.05$. Bar: Standard deviation.

N-SH and LA-N-5 (9). From the sequenced reads, those mapped to a reference genome sequence were selected for peak detection (in other word, for detection of candidates for *LMO1*-binding site) and applied to 6 peak-detection programs with 9 different settings. A definite peak was defined as the peaks detected by more than 5 of the 9 settings of the programs in each cell line (9). In this study, we focused on the definite peaks located in or near known microRNA genes (within a 5-kb proximity of known genes), common to the two cell lines, and identified 5 microRNA genes; *hsa-mir-548ap*, *hsa-mir-198*, *hsa-mir-4461*, *hsa-mir-3648* and *hsa-mir-3687* (Table I). We examined the effect of *LMO1* suppression on these genes by performing microRNA microarray expression analyses, comparing the SK-N-SH NB cells in which *LMO1* was suppressed with shRNA, to its counterpart introduced with shRNA for a GFP gene as a reference (Figure 1). We listed the microRNAs that were down-regulated to half or less in NB cells following *LMO1* suppression, compared to reference (Table II). The *hsa-mir-3648* (Table I) was one of the listed microRNAs and its down-regulation was confirmed by real-time RT-PCR in both SK-N-SH and LA-N-5 cells where *LMO1* expression was suppressed (Figure 2). Assuming that *hsa-mir-3648* is an oncogenic microRNA up-regulated by *LMO1*, the consequence of inhibiting the microRNA in NB cells was examined. In addition to SK-N-SH and LA-N-5 cell lines, the SK-N-BE(2) cell line was used in the inhibition assay. The reason is that SK-N-SH and LA-N-5 have a low siRNA transfection efficiency and

SK-N-BE(2) is more permissive for siRNA transfection (Figure 3). An oligonucleotide inhibitor for *hsa-mir-3648* was introduced into the three NB cell lines and showed a very weak but statistically significant suppressive effect on LA-N-5, however, it did not suppress cell proliferation of the other two cell lines. This suggests that *hsa-mir-3648* may not be related to the cell-proliferation promoting activity of *LMO1* or may have just a limited role in the *LMO1* activity (Figure 4).

MicroRNAs of the let-7 family are indirectly regulated by LMO1. Next, we focused on microRNAs which were up-regulated in SK-N-SH cells, in which *LMO1* is suppressed. A total of 18 microRNAs were found to be up-regulated by 2- or more fold in SK-N-SH cells following suppression of *LMO1* (Table III). As none of the 18 genes is a regulatory target of *LMO1* identified by the ChIP-Seq analyses, all of them are likely to be under an indirect regulation by *LMO1*. Importantly, they included *hsa-miR-34a-5p* and *let-7* family microRNAs which were previously reported to suppress the NB oncogene MYCN (15-17). We examined the effects of the 8 microRNA, *hsa-miR-34a-5p*, *hsa-let-7a-5p*, *hsa-let-7b-5p*, *hsa-let-7c*, *hsa-let-7e-5p*, *hsa-let-7f-5p*, *hsa-let-7g-5p* and *hsa-let-7i-5p*, on cell proliferation by introducing their oligonucleotide mimic into NB cell lines. As a result, all the mimics showed a statistically significant inhibitory effect on the proliferation of, at least, one of the three cell lines (Figure 5). It is noteworthy that the inhibitory effect of the mimics

Table II. *MicroRNA down-regulated in SK-N-SH by LMO1 suppression.*

MicroRNA	shGFP		shLMO1		Log2 ratio	MicroRNA	shGFP		shLMO1		Log2 ratio
	Signal*	Call†	Signal*	Call†			Signal*	Call†	Signal*	Call†	
hsa-miR-6165	19.3	1	0.1	0	-7.59	hsa-miR-5703	224.3	1	76.1	1	-1.56
hsa-miR-483-5p	16.8	1	0.1	0	-7.39	hsa-miR-630	341.3	1	119.5	1	-1.51
hsa-miR-4433-3p	15.7	1	0.1	0	-7.30	hsa-miR-5001-5p	70.1	1	25.9	1	-1.44
hsa-miR-4800-5p	15.1	1	0.1	0	-7.24	hsa-miR-4281	567.4	1	212.2	1	-1.42
hsa-miR-4298	13.7	1	0.1	0	-7.10	hsa-miR-4763-3p	45.4	1	17.0	1	-1.42
hsa-miR-4669	13.0	1	0.1	0	-7.03	hsa-miR-1246	458.4	1	175.4	1	-1.39
hsa-miR-125a-3p	12.0	1	0.1	0	-6.91	hsa-miR-4466	91.7	1	35.1	1	-1.39
hsa-miR-4442	10.0	1	0.1	0	-6.65	hsa-miR-6124	38.0	1	14.7	1	-1.37
hsa-miR-4465	10.0	1	0.1	0	-6.64	hsa-miR-371b-5p	58.4	1	22.9	1	-1.35
hsa-miR-4515	8.9	1	0.1	0	-6.48	hsa-miR-574-5p	15.2	1	6.0	1	-1.34
hsa-miR-4778-5p	8.6	1	0.1	0	-6.43	hsa-miR-5100	13.8	1	5.6	1	-1.31
hsa-miR-937-5p	8.5	1	0.1	0	-6.42	hsa-miR-6088	218.6	1	88.3	1	-1.31
hsa-miR-3648	8.4	1	0.1	0	-6.38	hsa-miR-5787	724.7	1	296.6	1	-1.29
hsa-miR-5195-3p	8.2	1	0.1	0	-6.35	hsa-miR-371a-5p	13.0	1	5.4	1	-1.26
hsa-miR-4417	8.1	1	0.1	0	-6.33	hsa-miR-4532	30.3	1	12.7	1	-1.25
hsa-miR-424-3p	7.9	1	0.1	0	-6.30	hsa-miR-6090	649.2	1	284.7	1	-1.19
hsa-miR-4793-5p	7.1	1	0.1	0	-6.16	hsa-miR-3196	105.7	1	46.4	1	-1.19
hsa-miR-6722-3p	6.3	1	0.1	0	-5.98	hsa-miR-3656	60.0	1	26.6	1	-1.17
hsa-miR-1185-1-3p	5.9	1	0.1	0	-5.87	hsa-miR-4787-5p	79.4	1	35.3	1	-1.17
hsa-miR-4430	5.3	1	0.1	0	-5.73	hsa-miR-6087	1214.1	1	542.0	1	-1.16
hsa-miR-3663-3p	4.7	1	0.1	0	-5.55	hsa-miR-6125	590.3	1	263.8	1	-1.16
hsa-miR-1233-1-5p	4.6	1	0.1	0	-5.53	hsa-miR-1915-3p	220.3	1	98.7	1	-1.16
hsa-miR-6723-5p	4.1	1	0.1	0	-5.36	hsa-miR-2392	8.2	1	3.7	1	-1.14
hsa-miR-874	3.8	1	0.1	0	-5.26	hsa-miR-1227-5p	19.9	1	9.1	1	-1.13
hsa-miR-664b-5p	3.8	1	0.1	0	-5.26	hsa-miR-3135b	19.1	1	8.8	1	-1.12
hsa-miR-1249	3.6	1	0.1	0	-5.15	hsa-miR-940	11.8	1	5.4	1	-1.12
hsa-miR-513a-5p	3.4	1	0.1	0	-5.11	hsa-miR-1973	74.1	1	34.6	1	-1.10
hsa-miR-642b-3p	24.8	1	4.2	1	-2.56	hsa-miR-4530	386.5	1	182.0	1	-1.09
hsa-miR-6510-5p	40.1	1	7.4	1	-2.43	hsa-miR-4534	49.8	1	23.7	1	-1.07
hsa-miR-1268a	680.3	1	151.7	1	-2.17	hsa-miR-4516	4300.4	1	2066.1	1	-1.06
hsa-miR-134	16.2	1	3.7	1	-2.13	hsa-miR-3940-5p	18.7	1	9.0	1	-1.05
hsa-miR-4270	34.0	1	8.1	1	-2.06	hsa-miR-1202	116.2	1	56.2	1	-1.05
hsa-miR-1290	322.5	1	78.4	1	-2.04	hsa-miR-3665	120.2	1	58.7	1	-1.03
hsa-miR-4499	15.7	1	3.8	1	-2.04	hsa-miR-1229-5p	60.9	1	29.8	1	-1.03
hsa-miR-575	37.2	1	10.8	1	-1.78	hsa-miR-3679-5p	57.7	1	28.9395	1	-1.00
hsa-miR-6126	13.4	1	3.9	1	-1.77						
hsa-miR-939-5p	29.1	1	8.8	1	-1.72						
hsa-miR-4741	73.6	1	22.9	1	-1.68						
hsa-miR-663a	22.3	1	7.1	1	-1.65						
hsa-miR-4745-5p	18.0	1	5.8	1	-1.65						
hsa-miR-1234-5p	492.4	1	165.6	1	-1.57						

*Exp gTotalGeneSignal (This signal is the sum of the total probe signals in the green channel per gene). †1 for detection of expression of mature microRNA, 0 for no detection.

Discussion

was clearly demonstrated especially in SK-N-BE(2), which display a much higher transfection efficiency for siRNA than SK-N-SH or LA-N-5.

Finally, the effect of the other 10 of the 18 microRNAs on cell proliferation was examined. Mimics for three microRNAs, *hsa-miR-16-5p*, *hsa-miR-20b-5p* and *hsa-miR-409-3p* significantly suppressed cell growth of SK-N-BE(2) but not that of SK-N-SH or LA-N-5, which may reflect their differences in transfection efficiency (Figure 6).

In this study, *hsa-miR-3648* was identified as a direct target of LMO1, however, no cell proliferation inhibitory activity of the microRNA was demonstrated in NB cells. There is no previous report on the functions of *hsa-miR-3648*. A recent report stated that *hsa-miR-3648* is transcriptionally regulated in neural stem cells by the amyloid precursor protein (APP)-derived cleavage product, APP intracellular domain (AICD) (18). Taken together with our results indicating that *hsa-miR-3648* is directly regulated by LMO1, we can suggest that

Figure 2. Down-regulation of *hsa-miR-3648* by shRNA (real-time RT-PCR). The shRNA for LMO1 (shLMO1) down-regulates *hsa-miR-3648* expression in NB cell lines, SK-N-SH and LA-N-5, compared to the cells transfected with shRNA for green fluorescent protein (shGFP), statistical significance at $p < 0.05$. Bar: Standard deviation.

AICD may cooperatively regulate the expression of *hsa-miR-3648* with LMO1 in NB cells.

The *MYCN* gene is the best-known oncogene that functions in NB and promotes cell proliferation, cell invasion and angiogenesis by regulating a variety of molecules (17). In the *let-7* family, *let-7e* was demonstrated to bind to the 3' untranslated region of the *MYCN* transcript and inhibit its function, conferring a tumor suppressor activity in NB (15). Humans have 10 mature *let-7* family members; *let-7a*, *let-7b*, *let-7c*, *let-7d*, *let-7e*, *let-7f*, *let-7g*, *let-7i*, *mir-98* and *mir-202* (19). Also, a tumor suppressive function of some *let-7* family members was demonstrated in non-NB cancers. Stable transfection of *let-7f* reduced growth of thyroid cancer cells through inhibition of mitogen-activated protein kinase (20). It was reported that proliferation of lung cancer cells is suppressed by *let-7a*, *let-7b*, *let-7c*, *let-7d* and *let-7g* (21). Proliferation of nasopharyngeal carcinoma cell lines, HK1 and HONE1, was inhibited by *let-7a*, *let-7b*, *let-7d*, *let-7e*, *let-7g* and *let-7i* (22). By targeting MYH9 (Myosin, Heavy Chain 9, Non-Muscle) transcripts, *let-7f* inhibits invasion and metastasis of gastric cancer cells (23). A malignant phenotype of osteosarcoma cell lines, U2-OS and HOS, is changed by *let-7g* through inhibition of Aurora kinase B gene (24).

LIN28B is an RNA binding protein that binds to the terminal loop of *let-7* microRNAs and inhibits processing of the microRNA (25). As silencing of LIN28B in NB cells up-regulates expression of *let-7a*, *let-7e*, *let-7g*, *let-7i*, and *mir-98*, it is likely that these microRNAs are regulated by

Figure 3. SK-N-BE(2) is more permissive to siRNA transfection than SK-N-SH. A western blot analysis demonstrated that introduction of siRNA for GAPDH reduced the amount of the GAPDH in SK-N-BE(2) cells more than in SK-N-SH.

LIN28B (26). A genome wide association study revealed *LIN28B* as an NB susceptibility gene, and Lin28b-transgenic mice develop NB (26). Intriguingly, Lin28 is down-regulated by *let-7* and both molecules form a reciprocal autoregulatory loop that controls microRNA processing during embryonic neural stem cell commitment in mice (27).

In addition to the *let-7* family, miR-34a was reported to target *MYCN* transcripts (16). It was also reported as a tumor suppressor that inhibits proliferation of NB cells through

Figure 4. The microRNA hsa-mir-3648 has no cell-proliferation promoting activity (cell growth assay). Transfection of hsa-mir-3648 inhibitor into NB cell lines showed no suppression of cell proliferation compared to negative control mimic (Neg cont). p-Value between the negative control (Neg Cont) and the inhibitor is shown. Bar: Standard deviation.

targeting mitogen-activated protein kinase kinase 9 (MAP3K9) transcripts (28, 29).

Besides the 7 microRNAs of the *let-7* family and miR-34a, this study detected 10 microRNAs indirectly suppressed by LMO1. Among these, miR-20b, miR-16 and miR-409 were clearly demonstrated to have a cell proliferation activity on SK-N-BE(2) (Table III, Figure 6). However, their targets in normal adrenal tissue or NB remain to be elucidated. Up-regulation of miR-20b expression was reported in gastric, breast and colorectal cancers and it is thought to act as an oncomir (30-32). However, an inhibitory activity of miR-20b on proliferation, migration and invasion of bladder cancer cells was also demonstrated (33). It may have a role in both tumor progression and suppression in a cancer type-dependent manner. It was reported that cisplatin down-regulates the expression of brain-derived neurotropic factor, a member of the neurotrophin family associated with aggressive malignant behavior by targeting miR-16 and resulting in inhibition of cell proliferation of the NB cell line SH-SY5Y (34). miR-409 suppresses cancer cell invasion and metastasis by targeting a pro-metastatic gene radixin in gastric cancers (35).

The results of this study suggest that LMO1 executes its oncogenic function in NB, partly by down-regulating several tumor suppressive microRNAs.

Acknowledgements

This work was supported by the National Cancer Center Research and Development Fund (No. 23-B-18) and by JSPS KAKENHI Grant Number 26670510.

Table III. MicroRNA up-regulated in SK-N-SH by LMO1 suppression.

MicroRNA	shGFP		shLMO1		Log2 ratio
	Signal*	Call†	Signal*	Call†	
hsa-miR-34a-5p	0.1	0	7.04682	1	6.14
hsa-miR-23a-3p	0.1	0	5.77634	1	5.85
hsa-miR-99b-5p	0.1	0	5.23699	1	5.71
hsa-miR-20b-5p	0.1	0	5.05015	1	5.66
hsa-let-7i-5p	0.1	0	4.88249	1	5.61
hsa-miR-106b-5p	0.1	0	4.69304	1	5.55
hsa-miR-543	0.1	0	3.92212	1	5.29
hsa-let-7g-5p	3.98429	1	16.964	1	2.09
hsa-let-7c	16.1552	1	42.2622	1	1.39
hsa-let-7f-5p	27.0994	1	70.6598	1	1.38
hsa-let-7b-5p	15.5918	1	40.6471	1	1.38
hsa-miR-16-5p	7.9961	1	20.0237	1	1.32
hsa-miR-125b-5p	39.1198	1	92.9244	1	1.25
hsa-miR-342-3p	10.0142	1	23.3954	1	1.22
hsa-miR-125a-5p	7.70954	1	17.639	1	1.19
hsa-let-7a-5p	92.7249	1	210.23	1	1.18
hsa-let-7e-5p	27.6341	1	59.0392	1	1.10
hsa-miR-409-3p	5.24178	1	11.1176	1	1.08

*Exp gTotalGeneSignal (This signal is the sum of the total probe signals in the green channel per gene). †1 for detection of expression of mature microRNA, 0 for no detection.

References

- Maris JM, Hogarty MD, Bagatell R and Cohn SL: Neuroblastoma. *Lancet* 369: 2106-2120, 2007.
- Maris JM: Recent advances in neuroblastoma. *N Engl J Med* 362: 2202-2211, 2010.

Figure 5. Seven *let-7* family microRNA mimics and *hsa-miR-34a-5p* mimic suppress NB cell growth. Introduction of mimic of each microRNA into NB cell lines resulted in reduction of cell proliferation compared to negative control mimic (Neg cont). Asterisks indicate statistical significance (*p*-value less than 0.05) between the negative control (Neg cont) and the mimic. Bar: Standard deviation.

Figure 6. MicroRNA mimics for *hsa-miR-16-5p*, *hsa-miR-20b-5p* and *hsa-miR-409-3p* suppress cell growth of SK-N-BE(2). Introduction of mimics for the three microRNA reduced cell proliferation of SK-N-BE(2) compared to negative control mimic (Neg cont). The mimic for *hsa-miR-23a-3p* showed a weak cell proliferation inhibitory effect on SK-N-SH. Asterisks indicate statistical significance (*p*-value less than 0.05) between the negative control (Neg cont) and the mimic. Bar: Standard deviation.

- 3 Morgenstern DA, Baruchel S and Irwin MS: Current and future strategies for relapsed neuroblastoma: challenges on the road to precision therapy. *J Pediatr Hematol Oncol* 35: 337-347, 2013.
- 4 Song HY, Rellinger EJ, Park SH, Paul P, Qiao J, Vasilopoulos A, Ozden O, Gius D and Chung DH: Inhibition of Sirtuin 6 induces neuroblastoma differentiation. *Anticancer Res* 38: 647-654, 2018.
- 5 Berger M and VON Schweinitz D: Therapeutic innovations for targeting childhood neuroblastoma: Implications of the neurokinin-1 receptor system. *Anticancer Res* 37: 5911-5918, 2017.

- 6 Abbou S, Lanvers-Kaminsky C, Daudigeos-Dubus E, LE Dret L, Laplace-Builhe C, Molenaar J, Vassal G, Geoerger B; within the ITCC Biology and Preclinical Evaluation Committee: Polo-like kinase inhibitor volasertib exhibits antitumor activity and synergy with vincristine in pediatric malignancies. *Anticancer Res* 36: 599-609, 2016.
- 7 Wang K, Diskin SJ, Zhang H, Attiyeh EF, Winter C, Hou C, Schnepf RW, Diamond M, Bosse K, Mayes PA, Glessner J, Kim C, Frackelton E, Garriss M, Wang Q, Glaberson W, Chiavacci R, Nguyen L, Jagannathan J, Saeki N, Sasaki H, Grant SF, Iolascon A, Mosse YP, Cole KA, Li H, Devoto M, McGrady PW, London WB, Capasso M, Rahman N, Hakonarson H and Maris JM: Integrative genomics identifies LMO1 as a neuroblastoma oncogene. *Nature* 469: 216-220, 2011.

- 8 Oldridge DA, Wood AC, Weichert-Leahey N, Crimmins I, Sussman R, Winter C, McDaniel LD, Diamond M, Hart LS, Zhu S, Durbin AD, Abraham BJ, Anders L, Tian L, Zhang S, Wei JS, Khan J, Bramlett K, Rahman N, Capasso M, Iolascon A, Gerhard DS, Guidry Auvil JM, Young RA, Hakonarson H, Diskin SJ, Look AT and Maris JM: Genetic predisposition to neuroblastoma mediated by a LMO1 super-enhancer polymorphism. *Nature* 528: 418-421, 2015.
- 9 Saeki N, Saito A, Sugaya Y, Amemiya M, Ono H, Komatsuzaki R, Yanagihara K and Sasaki H: Chromatin immunoprecipitation and DNA sequencing identified a LIMS1/ILK pathway regulated by an oncogenic transcription factor LMO1 in neuroblastoma. *Cancer Genomics Proteomics* 15: 165-174, 2018.
- 10 Hannigan G, Troussard AA and Dedhar S: Integrin-linked kinase: a cancer therapeutic target unique among its ILK. *Nat Rev Cancer* 5: 51-63, 2005.
- 11 Lee SL, Hsu EC, Chou CC, Chuang HC, Bai LY, Kulp SK and Chen CS: Identification and characterization of a novel integrin-linked kinase inhibitor. *J Med Chem* 54: 6364-6374, 2011.
- 12 Dougherty GW, Jose C, Gimona M and Cutler ML: The Rsu-1-PINCH1-ILK complex is regulated by Ras activation in tumor cells. *Eur J Cell Biol* 87: 721-734, 2008.
- 13 Krol J, Loedige I and Filipowicz W: The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet* 11: 597-610, 2010.
- 14 Bartel DP: Metazoan MicroRNAs. *Cell* 173: 20-51, 2018.
- 15 Buechner J, Tømte E, Haug BH, Henriksen JR, Løkke C, Flægstad T and Einvik C: Tumour-suppressor microRNAs let-7 and mir-101 target the proto-oncogene MYCN and inhibit cell proliferation in MYCN-amplified neuroblastoma. *Br J Cancer* 105: 296-303, 2011.
- 16 Wei JS, Song YK, Durinck S, Chen QR, Cheuk AT, Tsang P, Zhang Q, Thiele CJ, Slack A, Shohet J and Khan J: The MYCN oncogene is a direct target of miR-34a. *Oncogene* 27: 5204-5213, 2008.
- 17 Gherardi S, Valli E, Enriquez D and Perini G: MYCN-mediated transcriptional repression in neuroblastoma: the other side of the coin. *Front Oncol* 3: 42, 2013.
- 18 Shu R, Wong W, Ma QH, Yang ZZ, Zhu H, Liu FJ, Wang P, Ma J, Yan S, Polo JM, Bernard CC, Stanton LW, Dawe GS and Xiao ZC: APP intracellular domain acts as a transcriptional regulator of miR-663 suppressing neuronal differentiation. *Cell Death Dis* 6: e1651, 2015.
- 19 Roush S and Slack FJ: The let-7 family of microRNAs. *Trends Cell Biol* 18: 505-516, 2008.
- 20 Ricarte-Filho JC, Fuziwara CS, Yamashita AS, Rezende E, da-Silva MJ and Kimura ET: Effects of let-7 microRNA on cell growth and differentiation of papillary thyroid cancer. *Transl Oncol* 2: 236-241, 2009.
- 21 Johnson CD, Esquela-Kerscher A, Stefani G, Byrom M, Kelnar K, Ovcharenko D, Wilson M, Wang X, Shelton J, Shingara J, Chin L, Brown D and Slack FJ: The let-7 microRNA represses cell proliferation pathways in human cells. *Cancer Res* 67: 7713-7722, 2007.
- 22 Wong TS, Man OY, Tsang CM, Tsao SW, Tsang RK, Chan JY, Ho WK, Wei WI and To VS: MicroRNA let-7 suppresses nasopharyngeal carcinoma cells proliferation through down-regulating c-Myc expression. *J Cancer Res Clin Oncol* 137: 415-422, 2011.
- 23 Liang S, He L, Zhao X, Miao Y, Gu Y, Guo C, Xue Z, Dou W, Hu F, Wu K, Nie Y and Fan D: MicroRNA let-7f inhibits tumor invasion and metastasis by targeting MYH9 in human gastric cancer. *PLoS One* 6: e18409, 2011.
- 24 Liu JM, Long XH, Zhang GM, Zhou Y, Chen XY, Huang SH, Liu ZL and Zhang ZH: Let-7g reverses malignant phenotype of osteosarcoma cells by targeting Aurora-B. *Int J Clin Exp Pathol* 7: 4596-4606, 2014.
- 25 Piskounova E, Viswanathan SR, Janas M, LaPierre RJ, Daley GQ, Sliz P and Gregory RI: Determinants of microRNA processing inhibition by the developmentally regulated RNA-binding protein Lin28. *J Biol Chem* 283: 21310-21314, 2008.
- 26 Molenaar JJ, Domingo-Fernández R, Ebus ME, Lindner S, Koster J, Drabek K, Mestdagh P, van Sluis P, Valentijn LJ, van Nes J, Broekmans M, Haneveld F, Volckmann R, Bray I, Heukamp L, Sprüssel A, Thor T, Kieckbusch K, Klein-Hitpass L, Fischer M, Vandesompele J, Schramm A, van Noesel MM, Varesio L, Speleman F, Eggert A, Stallings RL, Caron HN, Versteeg R and Schulte JH: LIN28B induces neuroblastoma and enhances MYCN levels via let-7 suppression. *Nat Genet* 44: 1199-1206, 2012.
- 27 Rybak A, Fuchs H, Smirnova L, Brandt C, Pohl EE, Nitsch R and Wulczyn FG: A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* 10: 987-993, 2008.
- 28 Cole KA, Attiyeh EF, Mosse YP, Laquaglia MJ, Diskin SJ, Brodeur GM and Maris JM: A functional screen identifies miR-34a as a candidate neuroblastoma tumor suppressor gene. *Mol Cancer Res* 6: 735-742, 2008.
- 29 Tivnan A, Tracey L, Buckley PG, Alcock LC, Davidoff AM and Stallings RL: MicroRNA-34a is a potent tumor suppressor molecule in vivo in neuroblastoma. *BMC Cancer* 11: 33, 2011.
- 30 Xue TM, Tao LD, Zhang M, Xu GC, Zhang J and Zhang PJ: miR-20b overexpression is predictive of poor prognosis in gastric cancer. *Onco Targets Ther* 8: 1871-1876, 2015.
- 31 Ahmad A, Ginnebaugh KR, Sethi S, Chen W, Ali R, Mittal S and Sarkar FH: miR-20b is up-regulated in brain metastases from primary breast cancers. *Oncotarget* 6: 12188-12195, 2015.
- 32 Zhu J, Chen L, Zou L, Yang P, Wu R, Mao Y, Zhou H, Li R, Wang K, Wang W, Hua D and Zhang X: MiR-20b, -21, and -130b inhibit PTEN expression resulting in B7-H1 over-expression in advanced colorectal cancer. *Hum Immunol* 75: 348-353, 2014.
- 33 Park SL, Cho TM, Won SY, Song JH, Noh DH, Kim WJ and Moon SK: MicroRNA-20b inhibits the proliferation, migration and invasion of bladder cancer EJ cells via the targeting of cell cycle regulation and Sp-1-mediated MMP-2 expression. *Oncol Rep* 34: 1605-1612, 2015.
- 34 Sun YX, Yang J, Wang PY, Li YJ, Xie SY and Sun RP: Cisplatin regulates SH-SY5Y cell growth through downregulation of BDNF via miR-16. *Oncol Rep* 30: 2343-2349, 2013.
- 35 Zheng B, Liang L, Huang S, Zha R, Liu L, Jia D, Tian Q, Wang Q, Wang C, Long Z, Zhou Y, Cao X, Du C, Shi Y and He X: MicroRNA-409 suppresses tumour cell invasion and metastasis by directly targeting radixin in gastric cancers. *Oncogene* 31: 4509-4516, 2012.

Received July 8, 2018

Revised July 19, 2018

Accepted July 23, 2018